


## Unit 13

## Acids and Bases

1. Which compound is an electrolyte?  
 (A)  $C_6H_{12}O_6$  (C)  $CH_3OH$   
 (B)  $CaCl_2$  (D)  $CCl_4$
2. According to the Arrhenius theory, when a base dissolves in water it produces  
 (A)  $CO_3^{2-}$  as the only negative ion in solution  
 (B)  $OH^-$  as the only negative ion in solution  
 (C)  $NH_4^+$  as the only positive ion in solution  
 (D)  $H^+$  as the only positive ion in solution
3. Which substance is an Arrhenius acid?  
 (A)  $NH_3$  (C)  $HC_2H_3O_2$   
 (B)  $KOH$  (D)  $CH_3OH$
4. The pH of a 0.1 M solution is 11. What is the concentration of  $H_3O^+$  ions, in moles per liter?  
 (A)  $1 \times 10^{-1}$   
 (B)  $1 \times 10^{-3}$   
 (C)  $1 \times 10^{-11}$   
 (D)  $1 \times 10^{-13}$
5. What is the pH of a 0.01 M solution of  $HNO_3$ ?  
 (A) 1 (C) 13  
 (B) 2 (D) 14
6. Which reaction represents the process of neutralization?  
 (A)  $Mg(s) + 2 HCl(aq) \rightarrow MgCl_2(aq) + H_2(g)$   
 (B)  $HCl(aq) + KOH(aq) \rightarrow KCl(aq) + H_2O(l)$   
 (C)  $Pb(NO_3)_2(aq) + CaCl_2(aq) \rightarrow Ca(NO_3)_2(aq) + PbCl_2(s)$   
 (D)  $2 KClO_3(s) \rightarrow 2 KCl(s) + 3 O_2(g)$
7. Which acid-base pair will always undergo a reaction that produces a neutral solution?  
 (A) a weak acid and a weak base  
 (B) a weak acid and a strong base  
 (C) a strong acid and a weak base  
 (D) a strong acid and a strong base

8. Given the neutralization reaction:



Which compound is a salt?

- (A)  $KOH$  (C)  $K_2SO_4$   
 (B)  $H_2SO_4$  (D)  $H_2O$

9. An aqueous solution of an ionic compound turns red litmus blue, conducts electricity, and reacts with an acid to form a salt and water. This compound could be

- (A)  $HCl$  (C)  $KNO_3$   
 (B)  $NaI$  (D)  $LiOH$

10. When hydrochloric acid is neutralized by sodium hydroxide, the salt formed is sodium

- (A) hydrochlorate (C) chloride  
 (B) chlorate (D) perchloride

11. Which pH indicates a basic solution?

- (A) 1 (C) 7  
 (B) 5 (D) 12

12. Which of these pH numbers indicates the highest level of acidity?

- (A) 5 (C) 10  
 (B) 8 (D) 12

13. Which statement describes the characteristics of an Arrhenius base?

- (A) It changes blue litmus to red and has a pH less than 7.  
 (B) It changes blue litmus to red and has a pH greater than 7.  
 (C) It changes red litmus to blue and has a pH less than 7.  
 (D) It changes red litmus to blue and has a pH greater than 7.

14. Red litmus will turn blue when placed in an aqueous solution of

- (A)  $KCl$  (C)  $CH_3OH$   
 (B)  $KOH$  (D)  $CH_3COOH$

15. What is the  $H_3O^+$  ion concentration of a solution that has an  $OH^-$  ion concentration of  $1.0 \times 10^{-3} M$ ?

- (A)  $1.0 \times 10^{-3} M$   
 (B)  $1.0 \times 10^{-7} M$   
 (C)  $1.0 \times 10^{-11} M$   
 (D)  $1.0 \times 10^{-14} M$

16. What is the  $H^+$  ion concentration of an aqueous solution in which the  $OH^-$  ion concentration is  $1 \times 10^{-2}$  mole per liter?

- (A)  $1 \times 10^{-14} M$   
 (B)  $1 \times 10^{-12} M$   
 (C)  $1 \times 10^{-9} M$   
 (D)  $1 \times 10^{-2} M$

17. If a solution has a hydronium ion concentration of  $1 \times 10^{-9} M$ , the solution is

- (A) basic and has a pH of 9  
 (B) basic and has a pH of 5  
 (C) acidic and has a pH of 9  
 (D) acidic and has a pH of 5

18. Both  $HNO_3(aq)$  and  $CH_3COOH(aq)$  can be classified as

- (A) Arrhenius acids that turn blue litmus red  
 (B) Arrhenius bases that turn blue litmus red  
 (C) Arrhenius acids that turn red litmus blue  
 (D) Arrhenius bases that turn red litmus blue

19. A student records the following observations about an unknown solution:

- conducts electricity
- turns blue litmus red

The student should conclude that the unknown solution is most likely


- (A) an acid (C) an ester  
 (B) a base (D) an alcohol

20. Which acid is almost completely ionized in a dilute solution at 298K?

- (A)  $CH_3COOH$  (C)  $H_3PO_4$   
 (B)  $H_2S$  (D)  $HNO_3$



10. Which graph 1 presents the pressure-volume relationship for an ideal gas at constant temperature?



11. A gas sample has a volume of 25.0 milliliters at a pressure of 1.00 atmosphere. If the volume increases to 50.0 milliliters and the temperature remains constant, the new pressure will be

(A) 1.00 atm  
(B) 2.00 atm  
(C) 0.250 atm  
(D) 0.500 atm

16. When the pressure exerted on a confined gas at constant temperature is doubled, the volume of the gas is

(A) halved  
(B) doubled  
(C) tripled  
(D) quartered

17. If 4.00 moles of oxygen gas, 3.00 moles of hydrogen gas, and 1.00 mole of nitrogen gas are combined in a closed container at standard pressure, what is the partial pressure exerted by the hydrogen gas?

(A) 1.00 atm  
(B) 0.125 atm  
(C) 3.00 atm  
(D) 0.375 atm

18. A gas sample consisting of 2.0 moles of hydrogen and 1.0 mole of oxygen is collected over water at 29°C and 75 kPa. What is the partial pressure of the hydrogen in the sample?


(A) 24 kPa  
(B) 48 kPa  
(C) 72 kPa  
(D) 75 kPa

$T_{total} = 75 kPa = 3 atm$

$1.0 mol O_2 = 25 kPa$

18.2

15.




At constant pressure, which curve best shows the relationship between the volume of an ideal gas and its absolute temperature?

(A) A  
(B) B

(C) C  
(D) D

19. Which graph shows the pressure-temperature relationship expected for an ideal gas?



20. Real gas behavior deviates from ideal gas behavior because real gas particles have


(A) no volume and no attraction for each other  
(B) no volume but some attraction for each other  
(C) volume but no attraction for each other  
(D) volume and some attraction for each other

$T = avg KE$   
 $KE = \frac{1}{2} mv^2$

1. The average kinetic energy of water molecules is greatest in which of these samples?

(A) 10 g of water at 35°C  
(B) 10 g of water at 55°C  
(C) 100 g of water at 25°C  
(D) 100 g of water at 45°C

even the balanced equation representing a reaction:



Which statement is true about energy in this reaction?

(A) The reaction is exothermic because it releases heat.  
(B) The reaction is exothermic because it absorbs heat.  
(C) The reaction is endothermic because it releases heat.  
(D) The reaction is endothermic because it absorbs heat.

When ammonium chloride crystals are dissolved in water, the temperature of the water decreases. What does this temperature change indicate about the dissolving of ammonium chloride in water?

(A) It is an endothermic reaction because it absorbs heat.  
(B) It is an endothermic reaction because it releases heat.  
(C) It is an exothermic reaction because it absorbs heat.  
(D) It is an exothermic reaction because it releases heat.

4. As the temperature of a substance decreases, the average kinetic energy of its particles

(A) decreases  
(B) increases  
(C) remains the same  
(D) increases

5. When 200 grams of water cools from 50.°C to 25.°C, the total amount of heat energy released by the water is

(A) 42 kJ  
(B) 21 kJ  
(C) 34 J  
(D) 17 J

6. How many Joules of heat energy are released when 50. grams of water are cooled from 70.°C to 60.°C?

(A) 42 J  
(B) 210 J  
(C) 2100 J  
(D) 4200 J

7. A sample of water is heated from 10.0°C to 15.0°C by the addition of 126 Joules of heat. What is the mass of the water?

(A) 5.00 g  
(B) 6.00 g  
(C) 30.0 g  
(D) 150.0 g

$126 J = m \times 4.18 J/K \times 5^\circ C$

Unit 11

$\Delta T = 10^\circ C$

Thermochet


8. When 420 Joules of heat energy is added to 10. grams of water at 20.°C, the final temperature of the water will be

(A) 10.°C  
(B) 30.°C  
(C) 40.°C  
(D) 100°C

9. What is the total number of kilojoules of heat energy absorbed when the temperature of 200 grams of water is raised from 10°C to 40°C?

(A) 0.126 kJ  
(B) 0.840 kJ  
(C) 25.2 kJ  
(D) 33.6 kJ

10. The potential energy diagram below represents a reaction.



Which arrow represents the activation energy of the forward reaction?

(A) A  
(B) B  
(C) C  
(D) D

11. How much energy is required to vaporize 1000 grams of water at its boiling point?

(A) 2.26 kJ  
(B) 3.34 kJ  
(C) 4.2 kJ  
(D) 22.6 kJ

12. Approximately how many Joules of heat are needed to completely change 10.0 grams of ice to water at the melting point temperature?

(A) 1.00 J  
(B) 33.4 J  
(C) 334 J  
(D) 3,340 J


13. What is the total number of Joules lost when 10. grams of water at 80.°C is cooled to 60.°C?

(A) 42  
(B) 84  
(C) 420  
(D) 840



$420 J = 10 g \times 4.18 J/K \times \Delta T$

$42 = 10 \times 4.18 \times 20^\circ C$

14. The graph below represents the heating curve of a substance that starts as a solid below its freezing point.




- What is the melting point of this substance?  
(A) 30°C (B) 55°C
15. Base your answer to the following question on the potential energy diagram below, which represents the reaction:



- Which statement correctly describes this reaction?  
(A) It is endothermic and energy is absorbed.  
(B) It is endothermic and energy is released.  
(C) It is exothermic and energy is absorbed.  
(D) It is exothermic and energy is released.

- (C) 90°C (D) 120°C

16. Base your answer to the following question on the graph below. The graph shows heat being added at a constant rate to substance A and to substance B, which begin as solids below their melting point temperatures.



Compared to substance B, substance A has a


- (A) lower melting point and a lower boiling point  
(B) lower melting point and a higher boiling point  
(C) higher melting point and a lower boiling point  
(D) higher melting point and a higher boiling point

17. A 32 gram sample of iron (initially at 500 °C) is placed in 125 grams of water at 25°C, the final temperature of the system is 37.5°C. What is the specific heat of iron?

- (A) 42.2 (C) 45.7  
(B) 84.3 (D) 84.6

$$Q_m = 32g \times 4.18 J/g^\circ C \times 350$$

18. The graph below represents the uniform heating of a substance, starting with the substance as a solid below its melting point.



Which segment of the graph represents a time when both the solid and liquid phases are present?

- (A) AB (C) DE  
(B) BC (D) EF

19. How many Joules of heat energy are absorbed in raising the temperature of 10. grams of water from 5.0°C to 20.°C?

- (A) 3,959 J (C) 4,453 J  
(B) 39,590 J (D) 44,530 J

$$Q = 10g \times 4.18 J/g^\circ C \times 15$$

20. How many Joules of heat energy are required in raising the temperature of 100 grams of water from -10.0°C to 0°C?

- (A) -2,050 J (C) +2,050  
(B) -3,400 J (D) +3,400 J

$$Q = 100g \times 2.05 J/g^\circ C \times 10^\circ C = 2050 J$$

$$Q = 100g \times 2.05 J/g^\circ C \times 10^\circ C = 2050 J$$

$$500 - 375$$

$$Q = 32g \times 4.18 J/g^\circ C \times 350$$





- $(411 \text{ mL}) (790 \text{ kPa}) = (1880 \text{ mL}) (38.7 \text{ kPa})$   
 $T_2$   
 $273 \text{ K}$   
 work
1. A gas occupies a volume of 444 mL at 273 K and 79.0 kPa. What is the final kelvin temperature when the volume of the gas is changed to 1880 mL and the pressure is changed to 38.7 kPa?  
 (A) 31.5 K (C) 566 K  
 (B) 292 K (D) 2360 K

- $\frac{4.00 \text{ L}}{293 \text{ K}} = \frac{3.00 \text{ L}}{T_2}$
2. The volume of a gas is 4.00 liters at 293 K and constant pressure. For the volume of the gas to become 3.00 liters, the Kelvin temperature must be equal to  
 (A)  $3.00 \times 293$   
 4.00  
 (B)  $4.00 \times 293$   
 3.00  
 (C)  $3.00 \times 4.00$   
 293  
 (D)  $293$   
 3.00  $\times$  4.00

- $\frac{40.0 \text{ mL}}{293 \text{ K}} = \frac{80.0 \text{ mL}}{T_2}$   
 $20^\circ\text{C} \times \frac{80.0 \text{ mL}}{40.0 \text{ mL}} = 293 \text{ K} \times \frac{80.0 \text{ mL}}{40.0 \text{ mL}}$   
 (A) (C)  
 $20^\circ\text{C} \times \frac{40.0 \text{ mL}}{80.0 \text{ mL}} = 293 \text{ K} \times \frac{40.0 \text{ mL}}{80.0 \text{ mL}}$   
 (B) (D)

4. As the temperature of a gas increases at constant pressure, the volume of the gas  
 (A) decreases (C) remains the same  
 (B) increases Direct
5. Standard temperature and a pressure of 0.5 atmosphere are equal to  
 (A)  $0^\circ\text{C}$  and 51.6 kPa (C)  $0^\circ\text{C}$  and 101.3 kPa  
 (B)  $32^\circ\text{C}$  and 51.6 kPa (D)  $32^\circ\text{C}$  and 101.3 kPa

6. Which graph best shows the relationship between the pressure of a gas and its average kinetic energy at constant volume?



7. The temperature of a 2.0-liter sample of helium gas at STP is increased to  $27^\circ\text{C}$  and the pressure is decreased to 80. kPa. What is the new volume of the helium sample?  
 (A) 1.4 L (C) 2.8 L  
 (B) 2.0 L (D) 4.0 L

$(2.0 \text{ L}) (101.3 \text{ kPa}) = \frac{V_2 (80 \text{ kPa})}{300 \text{ K}}$   
 $(273 \text{ K})$

8. A gas has a volume of 1,400 milliliters at a temperature of 20. K and a pressure of 1.0 atm. What will be the new volume when the temperature is changed to 40. K and the pressure is changed to 0.50 atm?  
 (A) 350 mL (C) 1,400 mL  
 (B) 750 mL (D) 5,600 mL

9. A cylinder with a tightly fitted piston is shown in the diagram below.



Air at constant temperature

- As the piston moves downward, the number of molecules of air in the cylinder  
 (A) decreases (C) remains the same  
 (B) increases

12. The volume of a 1.00-mole sample of an ideal gas will decrease when the

- (A) pressure decreases and the temperature decreases  
 (B) pressure decreases and the temperature increases  
 (C) pressure increases and the temperature decreases  
 (D) pressure increases and the temperature increases

13. As the pressure of a gas at 150 kPa is changed to 100 kPa at constant temperature, the volume of the gas

- (A) decreases (C) remains the same  
 (B) increases

14. When 500. milliliters of hydrogen gas is heated from  $30^\circ\text{C}$  to  $60^\circ\text{C}$  at constant pressure, the volume of the gas at  $60^\circ\text{C}$  is equal to

$500 \text{ mL} \times \frac{213}{243}$   
 (A)



$500 \text{ mL} \times \frac{243}{213}$   
 (B)

$500 \text{ mL} \times \frac{333}{303}$   
 (C)

$500 \text{ mL} \times \frac{333}{273}$   
 (D)

$(500 \text{ mL}) \times \frac{V_2}{(273 \text{ K})} = \frac{V_2 (333 \text{ K})}{(273 \text{ K})}$


$\frac{273}{30} = \frac{273}{333}$



Unit 9

Stoichiometry


1. Given the equation:



What is the minimum number of liters of  $\text{CO}_2(\text{g})$ , measured at STP, needed to produce 32.0 grams of oxygen?

- A) 264 L C) 192 L  
B) 32.0 L D) 22.4 L


2. Given the unbalanced equation:



What is the coefficient of  $\text{O}_2$  when the equation is balanced correctly using the *smallest* whole number coefficients?

- A) 1 C) 3  
B) 2 D) 4


3. Given the reaction:



What is the total number of moles of water needed to make 2.5 moles of  $\text{C}_6\text{H}_{12}\text{O}_6$ ?

- A) 12 C) 2.5  
B) 6.0 D) 15


4. Given the balanced equation:



What is the total number of moles of  $\text{O}_2(\text{g})$  that must react completely with 5.00 moles of  $\text{C}_4\text{H}_{10}(\text{g})$ ?

- A) 32.5 C) 26.5  
B) 20.0 D) 10.0


5. Given the reaction:



What is the total number of grams of  $\text{O}_2(\text{g})$  needed to react completely with 0.50 mole of  $\text{C}_2\text{H}_2(\text{g})$ ?

- A) 160 g C) 40. g  
B) 80. g D) 10. g


6. Given the reaction



How many liters of ammonia, measured at STP, are produced when 28.0 grams of nitrogen is completely consumed?

- A) 44.8 C) 11.2  
B) 5.60 D) 22.4


7. Given the reaction:



What volume of  $\text{C}_8\text{H}_{18}(\text{g})$  will completely react to produce exactly 36 liters of  $\text{H}_2\text{O}(\text{g})$ ?

- A) 27 L C) 36 L  
B) 2.0 L D) 4.0 L


8. Given the equation:



How many moles of carbon dioxide are produced for each mole of butane consumed?

- A) 1 C) 8  
B) 2 D) 4


9. Given the reaction:



What is the total number of liters of  $\text{O}_2(\text{g})$  at STP needed to produce  $6.0 \times 10^{23}$  molecules of  $\text{H}_2\text{O}(\text{l})$ ?

- A) 33.6 L C) 44.8 L  
B) 22.4 L D) 11.2 L




10. Given the reaction:



The total number of grams of  $\text{O}_2$  needed to produce 54 grams of water is

- A) 84 C) 36  
B) 48 D) 75

P25






1. Which formula correctly represents antimony (V) oxide?

(A)  $\text{SbO}_5$  (C)  $\text{Sb}_2\text{O}_5$   
(B)  $\text{Sb}_3\text{O}$  (D)  $\text{Sb}_5\text{O}_2$

2. Given the reaction:



Which type of reaction is represented?

(A) single replacement (C) synthesis  
(B) double replacement (D) decomposition

3. Which equation represents a double replacement reaction?

(A)  $2 \text{Na} + 2 \text{H}_2\text{O} \rightarrow 2 \text{NaOH} + \text{H}_2$   
(B)  $\text{CaCO}_3 \rightarrow \text{CaO} + \text{CO}_2$   
(C)  $\text{LiOH} + \text{HCl} \rightarrow \text{LiCl} + \text{H}_2\text{O}$   
(D)  $\text{CH}_4 + 2 \text{O}_2 \rightarrow \text{CO}_2 + 2 \text{H}_2\text{O}$

4.  $2 \text{NH}_3(\text{g}) \leftrightarrow \text{N}_2(\text{g}) + 3 \text{H}_2(\text{g})$

What type of reaction is shown above?

(A) synthesis (C) single replacement  
(B) decomposition (D) double replacement

5.  $2 \text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \leftrightarrow 2 \text{SO}_3(\text{g})$

What type of reaction is shown above?

(A) synthesis (C) single replacement  
(B) decomposition (D) double replacement


6. When hydrocarbons burn completely in an excess of oxygen, the products are

(A) carbon monoxide and water  
(B) carbon dioxide and water  
(C) carbon monoxide and carbon dioxide  
(D) carbon dioxide and carbon

7. If an equation is balanced properly, both sides of the equation must have the same number of

(A) atoms (C) molecules  
(B) coefficients (D) moles of molecules


8. Given the unbalanced equation:



When the equation is balanced using the smallest whole-number coefficients, what is the coefficient of Al?

(A) 1 (C) 3  
(B) 2 (D) 4


9. Given the unbalanced equation:



When this equation is correctly balanced using smallest whole numbers, what is the coefficient of  $\text{O}_2(\text{g})$ ?

(A) 6 (C) 3  
(B) 2 (D) 4


10. Given the unbalanced equation:



What is the coefficient of  $\text{Al}_2(\text{SO}_4)_3$  when the equation is completely balanced using the smallest whole-number coefficients?

(A) 1 (C) 3  
(B) 2 (D) 4

11. When the equation




is balanced using smallest whole numbers, what is the coefficient of the  $\text{O}_2$ ?

(A) 1 (C) 3  
(B) 2 (D) 4

12. Which equation is correctly balanced?

(A)  $\text{CaO} + 2 \text{H}_2\text{O} \rightarrow \text{Ca(OH)}_2$   
(B)  $\text{NH}_3 + 2 \text{O}_2 \rightarrow \text{HNO}_3 + \text{H}_2\text{O}$   
(C)  $\text{Ca(OH)}_2 + 2 \text{H}_3\text{PO}_4 \rightarrow \text{Ca}_3(\text{PO}_4)_2 + 3 \text{H}_2\text{O}$   
(D)  $\text{Cu} + \text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 + \text{H}_2\text{O} + \text{SO}_2$


13. Given the unbalanced equation:



When the equation is completely balanced using the smallest whole number coefficients the sum of the coefficients is

(A) 5 (C) 3  
(B) 9 (D) 4


14. Given the unbalanced equation:



What is the coefficient in front of the  $\text{H}_2\text{O}$  when the equation is completely balanced using the smallest whole number coefficients?

(A) 6 (C) 3  
(B) 2 (D) 4


15. Given the balanced equation:



What is the correct formula for the product represented by the letter X?

(A)  $\text{MgO}$  (C)  $\text{MgO}_2$   
(B)  $\text{Mg}_2\text{O}$  (D)  $\text{Mg}_2\text{OH}$

16. Given the balanced equation:



What is the correct formula for the product represented by the letter X?


(A) K (C)  $\text{KCO}_3$   
(B) Cl (D) KCl

17. Given the incomplete equation:



Which set of products completes and balances the incomplete equation?

(A)  $\text{Ca} + \text{Cl}$  (C)  $\text{CaCl} + \text{O}_2$   
(B)  $\text{Ca} + \text{Cl}_2$  (D)  $\text{CaCl} + \text{H}_2\text{O}$



Which molecule is represented by X?

(A)  $\text{C}_2\text{H}_4$  (C)  $\text{C}_3\text{H}_6$   
(B)  $\text{C}_2\text{H}_6$  (D)  $\text{C}_3\text{H}_8$

1. What is the total number of moles of atoms present in 1 gram formula mass of  $\text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2$ ?  
 (A) 9 (B) 14 (C) 3 (D) 15
2. The gram formula mass of  $\text{NH}_4\text{Cl}$  is  
 (A) 22.4 g/mole (B) 28.0 g/mole (C) 53.5 g/mole (D) 95.5 g/mole
3. The gram-formula mass of  $(\text{NH}_4)_2\text{CO}_3$  is  
 (A) 46.0 g (B) 64.0 g (C) 78.0 g (D) 96.0 g
4. Which substance has the greatest molecular mass?  
 (A)  $\text{H}_2\text{O}_2$  (B) NO (C)  $\text{CF}_4$  (D)  $\text{I}_2$
5. The number of moles of molecules in a 12.0-gram sample of  $\text{Cl}_2$  is  
 (A)  $\frac{12.0}{35.5}$  mole (B)  $\frac{12.0}{71.0}$  mole (C) 12.0 moles (D)  $12.0 \times 35.5$  moles
6. The total number of moles represented by 20 grams of  $\text{CaCO}_3$  is  
 (A) 1 (B) 2 (C) 0.1 (D) 0.2
7. What is the total mass of 2.0 moles of  $\text{H}_2(\text{g})$ ?  
 (A) 1.0 g (B) 2.0 g (C) 3.0 g (D) 4.0 g
8. A sample of an unknown gas at STP has a density of 1.25 grams per liter. What is the gram molecular mass of this gas?  
 (A) 28.0 g (B) 44.0 g (C) 64.0 g (D) 80.0 g
9. The empirical formula of a compound is  $\text{CH}_3$ . The molecular formula of this compound could be  
 (A)  $\text{CH}_4$  (B)  $\text{C}_2\text{H}_4$  (C)  $\text{C}_2\text{H}_6$  (D)  $\text{C}_3\text{H}_6$
10. What is the empirical formula of a compound that contains 30.4% nitrogen and 69.6% oxygen by mass?  
 (A) NO (B)  $\text{NO}_2$  (C)  $\text{N}_2\text{O}_3$  (D)  $\text{N}_2\text{O}_5$
11. A compound consists of 25.9% nitrogen and 74.1% oxygen by mass. What is the empirical formula of the compound?  
 (A) NO (B)  $\text{NO}_2$  (C)  $\text{N}_2\text{O}$  (D)  $\text{N}_2\text{O}_5$
12. What is the percent by mass of oxygen in propanal,  $\text{CH}_3\text{CH}_2\text{CHO}$ ?  
 (A) 10.0% (B) 27.6% (C) 38.1% (D) 62.1%
13. In which compound is the percent by mass of oxygen greatest?  
 (A) BeO (B) MgO (C) CaO (D) SrO
14. An example of an empirical formula is  
 (A)  $\text{CH}_4$  (B)  $\text{C}_2\text{H}_4$  (C)  $\text{C}_2\text{H}_4(\text{OH})_2$  (D)  $\text{C}_6\text{H}_{12}\text{O}_6$
15. Which molecular formula is correctly paired with its corresponding empirical formula?  
 (A)  $\text{CO}_2$  and CO (B)  $\text{C}_2\text{H}_2$  and  $\text{CH}_2$  (C)  $\text{C}_6\text{H}_6$  and  $\text{C}_2\text{H}_2$  (D)  $\text{P}_4\text{O}_{10}$  and  $\text{P}_2\text{O}_5$

16. Which sample contains a total of  $9.0 \times 10^{23}$  atoms?  
 (A) 0.50 mole of HCl (B) 0.75 mole of  $\text{H}_2\text{O}$  (C) 1.5 moles of Cu (D) 1.5 moles of  $\text{H}_2$
17. What is the total number of atoms contained in a 1.00-mole sample of helium?  
 (A) 1.00 atom (B) 2.00 atoms (C)  $1.20 \times 10^{24}$  atoms (D)  $6.02 \times 10^{23}$  atoms
18. What is the total number of nitrogen atoms in 0.25 mole of  $\text{NO}_2$  gas?  
 (A)  $1.5 \times 10^{23}$  (B)  $6.0 \times 10^{23}$  (C)  $3.0 \times 10^{23}$  (D)  $1.2 \times 10^{24}$
19. The volume occupied by  $9.03 \times 10^{23}$  molecules of  $\text{N}_2$  gas at STP is closest to  
 (A) 0.500 liter (B) 1.50 liters (C) 22.4 liters (D) 33.6 liters
20. The total number of molecules in 34.0 grams of  $\text{NH}_3$  is equal to  
 (A)  $1.00 \times 22.4$  (B)  $2.00 \times 22.4$  (C)  $1.00 \times 6.02 \times 10^{23}$  (D)  $2.00 \times 6.02 \times 10^{23}$

$$\frac{12.0 \text{ g Cl}_2}{71.0 \text{ g/mol}} = 0.169 \text{ mol Cl}_2$$

$$\frac{20 \text{ g CaCO}_3}{100 \text{ g/mol}} = 0.2 \text{ mol CaCO}_3$$

$$\frac{1.25 \text{ g}}{1.01 \text{ g/L}} \times 22.4 \text{ L/mol} = 27.6 \text{ g/mol}$$

Smallest Molar mass

not E.F

correct

2 moles  
2 x  $\text{NH}_3$



1. The bonds between hydrogen and oxygen in a water molecule are classified as

(A) polar covalent (C) ionic  
(B) nonpolar covalent (D) metallic




2. Which molecule is nonpolar?

(A)  $H_2O$  (C)  $CO$   
(B)  $NH_3$  (D)  $CO_2$

3. Which of these substances has the strongest intermolecular forces?

(A)  $H_2O$  (C)  $H_2Se$   
(B)  $H_2S$  (D)  $H_2Te$

4. Which electron-dot structure represents a non-polar molecule?



5. Which molecule contains a triple covalent bond between its atoms?

(A)  $N_2$  (C)  $F_2$   
(B)  $O_2$  (D)  $H_2$

6. A diamond is an example of

(A) a supercooled liquid (C) a metallic substance  
(B) an ionic compound (D) a network solid

7. In which liquid is hydrogen bonding strongest?

(A)  $HF(l)$  (C)  $CH_3(l)$   
(B)  $H_2(l)$  (D)  $NH_3(l)$

8. Which type of molecule is  $CF_4$ ?

(A) polar, with a symmetrical distribution of charge  
(B) polar, with an asymmetrical distribution of charge  
(C) nonpolar, with a symmetrical distribution of charge  
(D) nonpolar, with an asymmetrical distribution of charge

9. The shape of a molecule of  $BF_3$  is said to be:

(A) trigonal planar (C) big bent  
(B) trigonal pyramidal (D) linear


10. Which compound has molecules that form the strongest hydrogen bonds?

(A)  $HI$  (C)  $HF$   
(B)  $HBr$  (D)  $HCl$

11. Which of the following compounds has the highest boiling point?

(A)  $H_2O$  (C)  $H_2Se$   
(B)  $H_2S$  (D)  $H_2Te$

12. Which pair of characteristics describes the molecule illustrated below?



2 lone pairs here

(A) symmetrical and polar  
(B) symmetrical and nonpolar  
(C) asymmetrical and polar  
(D) asymmetrical and nonpolar


13. Which molecule has an asymmetrical shape?

(A)  $N_2$  (C)  $Cl_2$   
(B)  $NH_3$  (D)  $CCl_4$


14. The shape of a molecule of  $BF_3$  is said to be:

(A) trigonal planar (C) big bent  
(B) trigonal pyramidal (D) linear

15. Which structural formula represents a nonpolar molecule?



16. Which electron-dot structure is correct for  $SiO_2$ ?



17. The shape of a molecule of oxygen is said to be:

(A) trigonal planar (C) big bent  
(B) trigonal pyramidal (D) linear

18. The shape of a molecule of  $PF_3$  is said to be:

(A) trigonal planar (C) big bent  
(B) trigonal pyramidal (D) linear

19. Which compound contains only covalent bonds?

(A)  $NaOH$  (C)  $Ca(OH)_2$   
(B)  $Ba(OH)_2$  (D)  $CH_3OH$

20. When phosphorus and chlorine atoms combine to form a molecule of  $PCl_3$ , 6 electrons will form

(A) nonpolar covalent bonds  
(B) polar covalent bonds  
(C) ionic bonds  
(D) hydrogen bonds

## Unit 5

## Bonding

1. Which formula represents an ionic compound?

- (A) NaCl (C) HCl  
(B) N<sub>2</sub>O (D) H<sub>2</sub>O

2. Which formula correctly represents the compound calcium hydroxide?

- (A) CaOH (C) CaOH<sub>2</sub>  
(B) Ca<sub>2</sub>OH (D) Ca(OH)<sub>2</sub>

3. Which metal will form a compound with the general formula M<sub>2</sub>CO<sub>3</sub> when it combines with a carbonate ion?

- (A) beryllium (C) calcium  
(B) aluminum (D) lithium

4. Which is the formula for magnesium sulfide?

- (A) MgS (C) MnS  
(B) MgSO<sub>3</sub> (D) MnSO<sub>3</sub>

5. The correct formula for calcium phosphate is

- (A) CaPO<sub>4</sub> (C) Ca<sub>3</sub>P<sub>2</sub>  
(B) Ca<sub>2</sub>(PO<sub>4</sub>)<sub>3</sub> (D) Ca<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>

6. What is the correct name of Fe<sub>2</sub>O<sub>3</sub>?

- (A) iron (I) oxide (C) iron (III) oxide  
(B) iron (II) oxide (D) iron (V) oxide

7. Which formula represents sodium sulfate?

- (A) NaSO<sub>4</sub> (C) Na<sub>2</sub>SO<sub>4</sub>  
(B) NaSO<sub>3</sub> (D) Na<sub>2</sub>SO<sub>3</sub>

8. Which formula correctly represents the composition of iron (III) oxide?

- (A) FeO<sub>3</sub> (C) Fe<sub>3</sub>O  
(B) Fe<sub>2</sub>O<sub>3</sub> (D) Fe<sub>3</sub>O<sub>2</sub>

9. If *M* represents a Group 1 metal, what is the formula for the compound formed by *M* and oxygen?

- (A) MO<sub>2</sub> (C) M<sub>2</sub>O<sub>3</sub>  
(B) M<sub>2</sub>O (D) M<sub>3</sub>O<sub>2</sub>

10. Element *M* is a metal and its chloride has the formula MCl<sub>2</sub>. To which group of the Periodic Table does element *M* most likely belong?

- (A) 1 (C) 15  
(B) 2 (D) 17

11. What is the correct name of the compound with the formula NH<sub>4</sub>NO<sub>2</sub>?

- (A) ammonia nitrite (C) ammonia nitrate  
(B) ammonium nitrite (D) ammonium nitrate

12. The chemical formula for nickel (II) bromide is

- (A) Ni<sub>2</sub>Br (C) N<sub>2</sub>Br  
(B) NiBr<sub>2</sub> (D) NBr<sub>2</sub>

13. Atoms of metals tend to

- (A) lose electrons and form negative ions  
(B) lose electrons and form positive ions  
(C) gain electrons and form negative ions  
(D) gain electrons and form positive ions

14. Which is the formula for the compound that forms when magnesium bonds with phosphorus?

- (A) Mg<sub>2</sub>P (C) Mg<sub>3</sub>P<sub>2</sub>  
(B) MgP<sub>2</sub> (D) Mg<sub>3</sub>P<sub>2</sub>

15. Which pair of atoms is held together by a covalent bond?

- (A) HCl (C) NaCl  
(B) LiCl (D) KCl

16. A correct name for N<sub>2</sub>O<sub>3</sub> is

- (A) nitrogen (I) oxide (C) nitrogen (III) oxide  
(B) nitrogen (II) oxide (D) nitrogen (IV) oxide

17. Which of the following is the correct formula for nitric acid?

- (A) HNO<sub>3</sub> (C) HF  
(B) HNO<sub>2</sub> (D) H<sub>2</sub>S

18. The name of the compound KClO<sub>2</sub> is potassium

- (A) hypochlorite (C) chlorate  
(B) chlorite (D) perchlorate

19. When a potassium atom reacts with bromine, the potassium atom will

- (A) lose only 1 electron (C) gain only 1 electron  
(B) lose 2 electrons (D) gain 2 electrons

20. What is the formula of nitrogen (II) oxide?

- (A) NO (C) N<sub>2</sub>O  
(B) NO<sub>2</sub> (D) N<sub>2</sub>O<sub>4</sub>

1. The element in Period 2 with the largest atomic radius is  
(A) a halogen (C) an alkali metal  
(B) a noble gas (D) an alkaline earth metal
2. Which sequence of atomic numbers represents elements which have similar chemical properties?  
(A) 19, 23, 30, 36 (C) 3, 12, 21, 40  
(B) 9, 16, 33, 50 (D) 4, 20, 38, 88
3. All of the atoms of the elements in Period 2 have the same number of  
(A) protons  
(B) neutrons  
(C) valence electrons  
(D) occupied energy levels (shells)
4. In which classification is an element placed if the outermost 3 sublevels of its atoms have a ground state electron configuration of  $3p^6 3d^5 4s^2$ ?  
(A) alkaline earth metals (C) metalloids (semimetals)  
(B) transition metals (D) nonmetals
5. Low ionization energies are most characteristic of atoms that are  
(A) metals (C) metalloids  
(B) nonmetals (D) noble gases
6. In a given period of the Periodic Table, the element with the lowest first ionization energy is always in  
(A) Group 1 (C) Group 17  
(B) Group 2 (D) Group 18
7. As the atoms of the elements in Group 1 are considered in order from top to bottom, compared to the ionization energy of the atom above it, the ionization energy of each successive atom  
(A) decreases (C) remains the same  
(B) increases
8. The table below shows some properties of elements A, B, C, and D.

Element	Ionization Energy	Electronegativity	Conductivity of Heat and Electricity
A	low	low	low
B	low	low	high
C	high	high	low
D	high	high	high

Which element is most likely a nonmetal?

- (A) A (C) C  
(B) B (D) D
9. Which of these metals loses electrons most readily?  
(A) calcium (C) potassium  
(B) magnesium (D) sodium
10. Which sequence correctly places the elements in order of increasing ionization energy?  
(A)  $H \rightarrow Li \rightarrow Na \rightarrow K$  (C)  $O \rightarrow S \rightarrow Se \rightarrow Te$   
(B)  $I \rightarrow Br \rightarrow Cl \rightarrow F$  (D)  $H \rightarrow Be \rightarrow Al \rightarrow Ga$
11. Which of the following particles has the smallest radius?  
(A)  $Na^0$   
(B)  $K^0$   
(C)  $Na^+$   
(D)  $K^+$

12. Which atom has the strongest attraction for electrons?

(A) Cl (C) Br  
(B) F (D) I

13. As the elements in Group 1 are considered in order of increasing atomic number, the atomic radius of each successive element increases. This is primarily due to an increase in the number of

(A) neutrons in the nucleus  
(B) electrons in the outermost shell  
(C) unpaired electrons  
(D) principal energy levels (shells)

14. When a sodium atom becomes an ion, the size of the atom

(A) decreases by gaining an electron  
(B) decreases by losing an electron  
(C) increases by gaining an electron  
(D) increases by losing an electron

15. Which element has an atomic radius that is greater than its ionic radius? metal

(A) S (C) F  
(B) K (D) O

16. Elements that readily gain electrons tend to have

(A) high ionization energy and high electronegativity  
(B) high ionization energy and low electronegativity  
(C) low ionization energy and low electronegativity  
(D) low ionization energy and high electronegativity


17. Which element in Period 3 has the greatest tendency to gain electrons?

(A) Na (C) Cl  
(B) Si (D) Ar

18. Which sequence of elements is arranged in order of decreasing atomic radii?

(A) Al, Si, P (C) Cl, Br, I  
(B) Li, Na, K (D) N, C, B

19. Which diagram correctly shows the relationship between electronegativity and atomic number for the elements of Period 3?



20. Within Period 2 of the Periodic Table, as the atomic number increases, the atomic radius generally

(A) decreases (C) remains the same  
(B) increases

1. Which particle has approximately the same mass as a proton?  
(A) alpha (C) electron  
(B) beta (D) neutron
2. Experimental evidence indicates that the nucleus of an atom  
(A) contains most of the mass of the atom  
(B) contains a small percentage of the mass of the atom  
(C) has no charge  
(D) has a negative charge
3. The atomic number of an atom is always equal to the total number of  
(A) neutrons in the nucleus  
(B) protons in the nucleus  
(C) neutrons plus protons in the atom  
(D) protons plus electrons in the atom
4. An atom that contains 35 protons, 45 neutrons, and 35 electrons has an atomic number of  
(A) 35 (C) 80  
(B) 45 (D) 115
5. Which atom has a nucleus that contains 13 protons and 14 neutrons?  
(A) Mg (C) Al  
(B) Be (D) N
6. What is the total number of electrons in a neutral atom of fluorine?  
(A) 9 (C) 19  
(B) 10 (D) 28
7. What is the mass number of an atom which contains 21 electrons, 21 protons, and 24 neutrons?  
(A) 21 (C) 45  
(B) 42 (D) 66
8. Which of the following particles has the least mass?  
(A) an electron (C) a hydrogen atom  
(B) a proton (D) a neutron
9. Compared to an atom of  $^{12}_6\text{C}$ , an atom of  $^{14}_6\text{C}$  has  
(A) more protons (C) more neutrons  
(B) fewer protons (D) fewer neutrons
10. Which two atoms are isotopes?  
(A)  $^{14}_6\text{C}$  and  $^{14}_7\text{N}$   
(B)  $^{23}_{11}\text{Na}$  and  $^{23}_{11}\text{Na}$   
(C)  $^3_1\text{H}$  and  $^3_2\text{He}$   
(D)  $^7_3\text{Li}$  and  $^9_4\text{Be}$
11. The number of protons in an atom of  $^3_1\text{H}$  is  
(A) 1 (C) 3  
(B) 2 (D) 4
12. The nucleus of an atom of  $^{127}_{53}\text{I}$  contains  
(A) 53 neutrons and 127 protons  
(B) 53 protons and 127 neutrons  
(C) 53 protons and 74 neutrons  
(D) 53 protons and 74 electrons
13. The nucleus of an atom of K-42 contains  
(A) 19 protons and 23 neutrons  
(B) 19 protons and 42 neutrons  
(C) 20 protons and 19 neutrons  
(D) 23 protons and 19 neutrons
14. An experiment in which alpha particles were used to bombard thin sheets of gold foil led to the conclusion that an atom is composed mostly of  
(A) empty space and has a small, negatively charged nucleus  
(B) empty space and has a small, positively charged nucleus  
(C) a large, dense, positively charged nucleus  
(D) a large, dense, negatively charged nucleus

15. Atoms of  $^{16}\text{O}$ ,  $^{17}\text{O}$ , and  $^{18}\text{O}$  have the same number of  
(A) neutrons, but a different number of protons  
(B) protons, but a different number of neutrons  
(C) protons, but a different number of electrons  
(D) electrons, but a different number of protons
16. All the isotopes of a given atom have  
(A) the same mass number and the same atomic number  
(B) the same mass number but different atomic numbers  
(C) different mass numbers but the same atomic number  
(D) different mass numbers and different atomic numbers
17. If the nucleus of an atom is represented as  $^{24}_{11}\text{X}$ , the atom is  
(A) Na (C) Mg  
(B) Al (D) Br
18. Which subatomic particles have a mass of approximately 1 atomic mass unit each?  
(A) proton and electron  
(B) proton and neutron  
(C) neutron and positron  
(D) electron and positron
19. Which statement concerning elements is true?  
(A) Different elements must have different numbers of isotopes.  
(B) Different elements must have different numbers of neutrons.  
(C) All atoms of a given element must have the same mass number.  
(D) All atoms of a given element must have the same atomic number.
20. The atomic mass of an element is defined as the weighted average mass of that element's  
(A) most abundant isotope  
(B) least abundant isotope  
(C) naturally occurring isotopes  
(D) radioactive isotopes